

Asymptotic behavior of positive solutions to a degenerate elliptic equation in the upper half space with a nonlinear boundary condition

Zhuoran Du

Abstract

We consider positive solutions of the problem

$$\begin{cases} -\operatorname{div}(x_n^a \nabla u) = 0 & \text{in } \mathbb{R}_+^n, \\ \frac{\partial u}{\partial \nu^a} = u^q & \text{on } \partial \mathbb{R}_+^n, \end{cases} \quad (1)$$

where $a \in (-1, 0) \cup (0, 1)$, $q > 1$ and $\frac{\partial u}{\partial \nu^a} := -\lim_{x_n \rightarrow 0^+} x_n^a \frac{\partial u}{\partial x_n}$. We obtain some qualitative properties of positive axially symmetric solutions in $n \geq 3$ for the case $a \in (-1, 0)$ under the condition $q \geq \frac{n-a}{n+a-2}$. In particular, we establish the asymptotic expansion of positive axially symmetric solutions.

Mathematics Subject Classification(2010): 35B07, 35B09, 35B40, 35J70.

Key words *Asymptotic behavior, positive solutions, degenerate elliptic equation, half space.*

1 Introduction

In this paper, we consider qualitative properties of positive solutions of the problem

$$\begin{cases} -\operatorname{div}(x_n^a \nabla u) = 0 & \text{in } \mathbb{R}_+^n = \{x \in \mathbb{R}^n : x_n > 0\}, \\ \frac{\partial u}{\partial \nu^a} = u^q & \text{on } \partial \mathbb{R}_+^n = \{x \in \mathbb{R}^n : x_n = 0\}, \end{cases} \quad (2)$$

where $a \in (-1, 0) \cup (0, 1)$, $q > 1$, $x = (x', x_n) \in \mathbb{R}^{n-1} \times [0, +\infty)$, and $\frac{\partial u}{\partial \nu^a} := -\lim_{x_n \rightarrow 0^+} x_n^a \frac{\partial u}{\partial x_n}$.

When the parameter $a = 0$, there had been many results about existence and qualitative properties of positive solutions to the problem (2). For critical case $q = \frac{n}{n-2}$, positive

⁰College of Mathematics and Econometrics, Hunan University, Changsha 410082, PRC. E-mail: duzr@hnu.edu.cn

solutions are known to exist in dimensions $n > 2$. Moreover, any positive solution is of the form (see [2] and [14])

$$u(x) = \alpha |x - x^0|^{-(n-2)}, \quad \alpha > 0, \quad x^0 = (x_1^0, \dots, x_n^0) \in \mathbb{R}^n, \quad x_n^0 = -\frac{1}{n-2} \alpha^{\frac{2}{n-2}}.$$

Positive solutions do not exist for subcritical case $q < \frac{n}{n-2}$ (see [12]). Subsequently, Chipot-Chlebk-Fila-Shafir [3] obtained the existence of positive solutions of (2) for the supercritical case $q > \frac{n}{n-2}$. Recently, under the conditions $n > 2$ and $q > \frac{n}{n-2}$, Harada [11] established the asymptotic expansion and the intersection property of positive solutions to the problem (2). Qui-Reichel [15] proved the existence of a unique singular positive axial symmetric solution for $q \geq \frac{n-1}{n-2}$, in $n \geq 3$.

It is known that the problem (2) with $a = 0$ is related to the square root of Laplacian equation

$$(-\Delta)^{\frac{1}{2}} v = v^q, \quad \text{in } \mathbb{R}^{n-1}. \quad (3)$$

Indeed, from the well-known Caffarelli-Silvestre extension in [1] we know that if u is a positive solution of (2), then a positive constant multiple of $u(x', 0) =: v(x')$ satisfies (3). Therefore the asymptotic expansion results in [11] generalize the asymptotic expansion results of the corresponding semilinear elliptic equation with standard Laplacian operator

$$-\Delta v = v^q, \quad \text{in } \mathbb{R}^{n-1}.$$

in [9], [13] to the square root of Laplacian operator case.

In the general case $a \in (-1, 0)$, Fang, Gui and the author in [5] obtain existence of positive axial symmetric solutions in space dimensions $n \geq 3$ for $q \geq \frac{n-a}{n+a-2}$. Further the author establish a unique singular positive axial symmetric solution $\psi_\infty(x)$, for $q \geq \frac{n-1}{n+a-2}$, in $n \geq 3$ for all $a \in (-1, 0) \cup (0, 1)$ (see [6]).

Naturally we hope to establish similar qualitative properties of the positive solution of (2) for the general case $a \in (-1, 0)$ as that for the case $a = 0$ in [11]. Our main results in this paper are stated as follows.

Theorem 1.1 *Let $q \geq \frac{n-a}{n+a-2}$ and denote $m_q = \frac{1-a}{q-1}$. Then there exists a family of positive axial symmetric solutions $u_\beta(x)$ ($u_\beta(0) = \beta > 0$) of (2) satisfying the following properties*

- (i) $u_\beta(x) = \beta u_1(\beta^{\frac{q-1}{1-a}} x)$;
- (ii) $u_\beta(x) \leq C(1 + |x|)^{-m_q}$;
- (iii) $\lim_{\beta \rightarrow \infty} u_\beta(x) = \psi_\infty(x)$ for $x \in \mathbb{R}_+^n \setminus \{0\}$, $q \neq \frac{n-a}{n+a-2}$.

The following Theorem is the asymptotic expansion for the JL-supercritical, JL-critical and JL-subcritical case (see Definition 3.1).

Theorem 1.2 (I) Let q be JL-supercritical or JL-critical. Then there exists $\varepsilon > 0$ such that for any $\beta > 0$ there exist $C_1(\beta) < 0, C_2(\beta) \in \mathbb{R}$, such that the solutions u_β of (2) hold the asymptotic expansion for large $r > 0$

$$u_\beta(x) = \psi_\infty(x) + [C_1 + O(r^{-\varepsilon})]e_1(\theta)r^{-(m_q+\rho_1)}, \quad \text{for JL-supercritical case,}$$

and

$$u_\beta(x) = \psi_\infty(x) + [C_1 \ln r + C_2 + O(r^{-\varepsilon})]e_1(\theta)r^{-\frac{n+a-2}{2}}, \quad \text{for JL-critical case,}$$

where we have used the polar coordinate $|x'| = r \sin \theta$, $x_n = r \cos \theta$, and

$$\rho_1 = \frac{(n+a-2-2m_q) - \sqrt{(n+a-2-2m_q)^2 + 4[m_q(n+a-2-m_q) + \lambda_1]}}{2},$$

where λ_i , $e_i(\theta)$ are the i -th eigenvalue, the i -th eigenfunction of (23) with

$$\int_0^{\frac{\pi}{2}} e_i(\theta)e_j(\theta) \sin^{n-2} \cos^a \theta d\theta = \delta_{ij}, \quad e_i\left(\frac{\pi}{2}\right) > 0.$$

Moreover the asymptotic expansion holds uniformly for $\theta \in (0, \frac{\pi}{2})$.

(II) Let q be JL-subcritical. Then the solutions u_β of (2) satisfies one of the following two asymptotic expansions for large $r > 0$

1) there exists $(c_1, c_2) \neq 0$ and $\varepsilon > 0$ such that

$$u_\beta(x) = \psi_\infty(x) + [c_1 \sin(K \ln r) + c_2 \cos(K \ln r) + O(r^{-\varepsilon})]e_1(\theta)r^{-\frac{n+a-2}{2}};$$

2) there exists $c > 0$ and $\varepsilon > 0$ such that

$$u_\beta(x) = \psi_\infty(x) + [c + O(r^{-\varepsilon})]e_2(\theta)r^{-(m_q+\rho_2)},$$

where $K > 0$, $\rho_2 > 0$ are given by

$$K = \frac{\sqrt{-(n+a-2-2m_q)^2 - 4[m_q(n+a-2-m_q) + \lambda_1]}}{2},$$

$$\rho_2 = \frac{(n+a-2-2m_q) + \sqrt{(n+a-2-2m_q)^2 + 4[m_q(n+a-2-m_q) + \lambda_2]}}{2}.$$

Moreover the asymptotic expansion holds uniformly for $\theta \in (0, \frac{\pi}{2})$.

Therefore, by using the Caffarelli-Silvestre extension, we know that our results generalize the results from the square root Laplacian operator to general fractional Laplacian operator

$$(-\Delta)^s v = v^q, \quad \text{in } \mathbb{R}^{n-1},$$

where $s = \frac{1-a}{2} \in (\frac{1}{2}, 1)$, since $a \in (-1, 0)$.

The paper is organized as follows. Section 2 contains some notations and an estimate of the infimum C_a in generalized trace Hardy inequality, which will be used in the definition of JL-critical exponent in Section 3. In section 4 estimates of eigenvalues of an eigenvalue problem are made, which will be used in asymptotic expansions for positive axial symmetric solutions. In section 5 we derive decay estimates and limit behavior of them. Section 6 contains the proof of asymptotic expansions for them.

2 Preliminaries

We say that a function u is axially symmetric with respect to the x_n -axis, if it can be expressed by $u(x) = \tilde{u}(|x'|, x_n)$ for some function $\tilde{u}$. For axially symmetric function, one need to introduce the polar coordinates (r, θ)

$$|x'| = r \sin \theta, \quad x_n = r \cos \theta, \quad \theta \in [0, \frac{\pi}{2}]. \quad (4)$$

We first recall the main results in [5] and [6].

Proposition 2.1 ([5]) *Let $q \geq \frac{n-a}{n+a-2}$ and $a \in (-1, 0)$. The problem (2) admits a positive axially symmetric solution $u(x)$ satisfying $u_\theta > 0$ and $u(0) = 1$.*

In [6], the author consider the existence of positive singular solutions of the form

$$\psi_\infty(x) = V(\theta)r^{\frac{a-1}{q-1}}.$$

To obtain $V(\theta)$, one need to solve

$$\begin{cases} (\sin^{n-2} \theta \cos^a \theta V_\theta)_\theta = \gamma V \sin^{n-2} \theta \cos^a \theta & \theta \in (0, \frac{\pi}{2}), \\ \lim_{\theta \rightarrow \frac{\pi}{2}} V_\theta(\theta) \cos^a \theta = V^q(\frac{\pi}{2}), \end{cases} \quad (5)$$

where

$$\gamma := m_q(n + a - 2 - m_q) > 0.$$

Proposition 2.2 ([6]) *Let $q \geq \frac{n-1}{n+a-2}$ and $a \in (-1, 0) \cup (0, 1)$. The problem (5) admits a unique solution $V(\theta)$ satisfying $V_\theta(\theta) > 0$.*

Next we introduce some notations that will be used. Let $S_+^{n-1} = \{\omega = (\omega', \cos \theta) : \omega' \in \mathbb{R}^{n-1}, \theta \in [0, \frac{\pi}{2}), |\omega'|^2 + \cos^2 \theta = 1\}$ be a half unit sphere. For $p > 1$, we define the space

$$L_{\text{sym}}^p(S_+^{n-1}) = \left\{ e : e(\omega) \text{ depends only on } \theta \text{ and } \int_0^{\frac{\pi}{2}} |e(\theta)|^p \sin^{n-2} \theta \cos^a \theta d\theta < +\infty \right\}.$$

We denote

$$\|e(\theta)\|_{p,a} := \left(\int_0^{\frac{\pi}{2}} |e(\theta)|^p \sin^{n-2} \theta \cos^a \theta d\theta \right)^{\frac{1}{p}}.$$

Further we define the space

$$\mathcal{H}_a^k(S_+^{n-1}) = \left\{ e \in L_{\text{sym}}^p(S_+^{n-1}) : \|e\|_{\mathcal{H}_a^k(S_+^{n-1})} := \sum_{i=0}^k \|\partial_\theta^i e(\theta)\|_{p,a} < +\infty \right\}.$$

We also define the following operator on S_+^{n-1} according to the norm $\|\cdot\|_{p,a}$

$$\Delta_{S,a} := \partial_{\theta\theta} + [(n-2)\cot\theta - a\tan\theta]\partial_{\theta}.$$

Note that

$$\sin^{n-2}\theta \cos^a\theta \Delta_{S,a}e(\theta) = (\sin^{n-2}\theta \cos^a\theta e_{\theta})_{\theta}.$$

We will establish the following elementary lemma.

Lemma 2.1 *For $a \in (-1, 0)$, we have*

$$\int_0^{\frac{\pi}{2}} \sin^{n+a-2}\theta d\theta < \int_0^{\frac{\pi}{2}} \sin^{n-2}\theta \cos^a\theta d\theta. \quad (6)$$

Proof. Since

$$\int_0^{\frac{\pi}{2}} \sin^{n+a-2}\theta d\theta = \int_0^{\frac{\pi}{4}} \sin^{n+a-2}\theta d\theta + \int_0^{\frac{\pi}{4}} \cos^{n+a-2}\theta d\theta$$

and

$$\int_0^{\frac{\pi}{2}} \sin^{n-2}\theta \cos^a\theta d\theta = \int_0^{\frac{\pi}{4}} \sin^{n-2}\theta \cos^a\theta d\theta + \int_0^{\frac{\pi}{4}} \cos^{n-2}\theta \sin^a\theta d\theta,$$

we have

$$\begin{aligned} & \int_0^{\frac{\pi}{2}} \sin^{n+a-2}\theta d\theta - \int_0^{\frac{\pi}{2}} \sin^{n-2}\theta \cos^a\theta d\theta \\ &= \int_0^{\frac{\pi}{4}} (\sin^{n-2}\theta - \cos^{n-2}\theta)(\sin^a\theta - \cos^a\theta) d\theta < 0. \end{aligned}$$

The lemma is proved.

We introduce the space $H_a^1(\mathbb{R}_+^n) := \left\{ u : \int_{\mathbb{R}_+^n} (u^2 + |\nabla u|^2) x_n^a dx < +\infty \right\}$, and consider the following minimizing problem

$$C_a := \inf_{u \in H_a^1(\mathbb{R}_+^n)} \frac{\int_{\mathbb{R}_+^n} |\nabla u|^2 x_n^a dx}{\int_{\partial\mathbb{R}_+^n} |u|^2 |x'|^{a-1} dx'},$$

which plays an essential role to define a so-called JL-critical exponent in Section 3.

Lemma 2.2 *Let $a \in (-1, 0)$ and $n \geq 4$. We have*

$$C_a > \frac{n+a-3}{2}.$$

Proof. One may assume that $u = u(\bar{r}, t)$ where $\bar{r} = |x'| = |(x_1, \dots, x_{n-1})|$ and $t = x_n$. Then

$$\int_{\partial \mathbb{R}_+^n} |u|^2 |x'|^{a-1} dx' = (n-1) \omega_{n-1} \int_0^\infty u^2(\bar{r}, 0) \bar{r}^{n+a-3} d\bar{r},$$

where ω_{n-1} denotes the measure of the unit sphere in $\mathbb{R}^{n-1}$. Since

$$u^2(\bar{r}, 0) = -2 \int_0^\infty u(\bar{r}, t) \partial_t u(\bar{r}, t) dt,$$

Hence

$$\begin{aligned} \int_{\partial \mathbb{R}_+^n} |u|^2 |x'|^{a-1} dx' &= -2(n-1) \omega_{n-1} \int_0^\infty \int_0^\infty u(\bar{r}, t) \partial_t u(\bar{r}, t) \bar{r}^{n+a-3} d\bar{r} dt \\ &\leq 2(n-1) \omega_{n-1} \int_0^\infty \left(\int_0^\infty u^2(\bar{r}, t) \bar{r}^{n+a-4} d\bar{r} \right)^{1/2} \left(\int_0^\infty (\partial_t u(\bar{r}, t))^2 \bar{r}^{n+a-2} d\bar{r} \right)^{1/2} dt. \end{aligned}$$

We use now the inequality

$$\int_0^\infty u^2(\bar{r}, t) \bar{r}^{n+a-4} d\bar{r} \leq \frac{4}{(n+a-3)^2} \int_0^\infty (\partial_{\bar{r}} u(\bar{r}, t))^2 \bar{r}^{n+a-2} d\bar{r},$$

which follows from the generalized Hardy's inequality

$$\int_{\mathbb{R}^d} \frac{u^2}{|x|^{2(b+1)}} dx \leq \frac{4}{(d-2-2b)^2} \int_{\mathbb{R}^d} \frac{|\nabla u|^2}{|x|^{2b}} dx.$$

We obtain

$$\begin{aligned} &\int_{\partial \mathbb{R}_+^n} |u|^2 |x'|^{a-1} dx' \\ &\leq \frac{4(n-1) \omega_{n-1}}{n+a-3} \int_0^\infty \left(\int_0^\infty (\partial_{\bar{r}} u(\bar{r}, t))^2 \bar{r}^{n+a-2} d\bar{r} \right)^{1/2} \left(\int_0^\infty (\partial_t u(\bar{r}, t))^2 \bar{r}^{n+a-2} d\bar{r} \right)^{1/2} dt \\ &\leq \frac{2(n-1) \omega_{n-1}}{n+a-3} \int_0^\infty \int_0^\infty [(\partial_{\bar{r}} u(\bar{r}, t))^2 + (\partial_t u(\bar{r}, t))^2] \bar{r}^{n+a-2} d\bar{r} dt. \end{aligned} \quad (7)$$

We introduce the coordinates (r, θ)

$$\bar{r} = r \sin \theta, \quad t = r \cos \theta.$$

Then one has

$$\begin{aligned} &\int_0^\infty \int_0^\infty [(\partial_{\bar{r}} u(\bar{r}, t))^2 + (\partial_t u(\bar{r}, t))^2] \bar{r}^{n+a-2} d\bar{r} dt \\ &= \int_0^\infty r^{n+a-1} \left(\int_0^{\frac{\pi}{2}} |\nabla u(r \sin \theta, r \cos \theta)|^2 \sin^{n+a-2} d\theta \right) dr. \end{aligned} \quad (8)$$

We may further assume that $|\nabla u(r \sin \theta, r \cos \theta)|^2 = |\nabla u(r \cos \theta, r \sin \theta)|^2$. From this, using the similar argument of Lemma 2.1, we obtain

$$\begin{aligned} & \int_0^\infty r^{n+a-1} \left(\int_0^{\frac{\pi}{2}} |\nabla u(r \sin \theta, r \cos \theta)|^2 \sin^{n+a-2} d\theta \right) dr \\ & < \int_0^\infty r^{n+a-1} \left(\int_0^{\frac{\pi}{2}} |\nabla u(r \sin \theta, r \cos \theta)|^2 \sin^{n-2} \cos^a \theta d\theta \right) dr. \end{aligned} \quad (9)$$

Note that

$$\begin{aligned} & \int_0^\infty r^{n+a-1} \left(\int_0^{\frac{\pi}{2}} |\nabla u(r \sin \theta, r \cos \theta)|^2 \sin^{n-2} \cos^a \theta d\theta \right) dr \\ & = \frac{1}{(n-1)\omega_{n-1}} \int_{\mathbb{R}_+^n} |\nabla u|^2 x_n^a dx. \end{aligned} \quad (10)$$

Hence, from (7)-(10), we have

$$\int_{\partial \mathbb{R}_+^n} |u|^2 |x'|^{a-1} dx' \leq \frac{2}{n+a-3} \int_{\mathbb{R}_+^n} |\nabla u|^2 x_n^a dx.$$

The proof of this lemma is complete.

Remark 1 This lemma generalize the conclusion that $C_0 > \frac{n-3}{2}$ for the case $a = 0$, which is obtained by Dávila-Dupaigne-Montenegro in [4].

We define

$$h_{n,a} := -\frac{(n+a-2)^2}{4}.$$

Next we show that $h_{n,a}$ is the first eigenvalue of the following eigenvalue problem

$$-\Delta_{S,a} e = \lambda e, \quad \theta \in (0, \frac{\pi}{2}), \quad \lim_{\theta \rightarrow \frac{\pi}{2}} e_\theta(\theta) \cos^a \theta = C_a e(\frac{\pi}{2}). \quad (11)$$

To this end, we introduce the space

$$H_a^1(S_+^{n-1}) := \left\{ e : \int_{S_+^{n-1}} (e^2 + |\nabla_S e|^2) d\omega = \int_{S_+^{n-1}} (e^2 + |\nabla_S e|^2) \sin^{n-2} \cos^a \theta d\theta d\omega' < +\infty \right\},$$

where $\nabla_S e$ is a gradient of e on S_+^{n-1} . We consider a minimizing problem

$$C_S := \inf_{e \in H_a^1(S_+^{n-1})} \frac{\|\nabla_S e\|_{L_a^2(S_+^{n-1})}^2 - h_{n,a} \|e\|_{L_a^2(S_+^{n-1})}^2}{\|e\|_{L^2(\partial S_+^{n-1})}^2}, \quad (12)$$

where $\|e\|_{L_a^2(S_+^{n-1})}^2 := \int_{S_+^{n-1}} e^2(\theta) \sin^{n-2} \cos^a \theta d\theta d\omega'$. For simplicity of notations, hereafter we set

$$d\mu := \sin^{n-2} \cos^a \theta d\theta, \quad e_B := e\left(\frac{\pi}{2}\right).$$

By a trace inequality, it is verified that there exists a minimizer $e \in \mathcal{H}_a^1(S_+^{n-1})$ of (12) satisfying $e(\theta) > 0$ and

$$-\Delta_{S,a} e = h_{n,a} e, \quad \theta \in (0, \frac{\pi}{2}), \quad \lim_{\theta \rightarrow \frac{\pi}{2}} e_\theta(\theta) \cos^a \theta = C_S e_B. \quad (13)$$

Since $e(\theta) > 0$, then $h_{n,a}$ is the first eigenvalue of this problem. Therefore, for problem (11), to prove $\lambda_1 = h_{n,a}$ we only need to show $C_S = C_a$.

Lemma 2.3 $C_S = C_a$.

Proof. Let $u \in C_c^1(\overline{\mathbb{R}_+^n})$. We have

$$\int_{\partial \mathbb{R}_+^n} |u(x', 0)|^2 |x'|^{a-1} dx' = \int_0^\infty \|u(r, \cdot)\|_{L^2(\partial S_+^{n-1})}^2 r^{n+a-3} dr.$$

By the definition of C_S and the generalized Hardy inequality, one has

$$\begin{aligned} C_S \int_0^\infty \|u(r, \cdot)\|_{L^2(\partial S_+^{n-1})}^2 r^{n+a-3} dr &\leq \int_0^\infty \int_{S_+^{n-1}} (|\nabla_S u|^2 - h_{n,a} u^2) r^{n+a-3} d\omega dr \\ &\leq \int_0^\infty \int_{S_+^{n-1}} (r^{-2} |\nabla_S u|^2 + u_r^2) r^{n+a-1} d\omega dr. \end{aligned}$$

Note that $|\nabla u|^2 = u_r^2 + r^{-2} |\nabla_S u|^2$, we obtain

$$C_S \int_0^\infty \|u(r, \cdot)\|_{L^2(\partial S_+^{n-1})}^2 r^{n+a-3} dr \leq \int_{\mathbb{R}_+^n} |\nabla u|^2 x_n^a dx,$$

which gives that $C_a \geq C_S$.

Next we show $C_a \leq C_S$. We need to construct a sequence $\{u_i\}_{i \in \mathbb{N}} \subset H_a^1(\mathbb{R}_+^n)$ such that

$$\lim_{i \rightarrow \infty} \frac{\int_{\mathbb{R}_+^n} |\nabla u_i|^2 x_n^a dx}{\int_{\partial \mathbb{R}_+^n} |u_i|^2 |x'|^{a-1} dx'} = C_S. \quad (14)$$

To this end, we set

$$u_i(x) = \begin{cases} e(\theta) i^{(n+a-2)/2} & \text{if } r \in [0, 1/i], \\ e(\theta) r^{(n+a-2)/2} & \text{if } r \in [1/i, 1), \\ \chi(r) e(\theta) r^{(n+a-2)/2} & \text{if } r \in [1, \infty), \end{cases} \quad (15)$$

where $e(\theta)$ is a minimizer of (12) and $\chi(r)$ is a cut-off function such that $\chi(r) = 1$ for $r \in [0, 1]$ and $\chi(r) = 0$ for $r \in [2, \infty)$. Calculation shows that

$$\int_{\mathbb{R}_+^n} |\nabla u_i|^2 x_n^a dx = C_S \|e\|_{L^2(\partial S_+^{n-1})}^2 (\ln i) + O(1)$$

and

$$\int_{\partial \mathbb{R}_+^n} |u(x', 0)|^2 |x'|^{a-1} dx' = \|e\|_{L^2(\partial S_+^{n-1})}^2 (\ln i) + O(1).$$

Hence we obtain (14). We complete the proof of this lemma.

3 JL-critical exponent

We introduce

$$J(q) := \inf_{u \in H_a^1(\mathbb{R}_+^n)} \frac{\int_{\mathbb{R}_+^n} |\nabla u|^2 x_n^a dx - q \int_{\partial \mathbb{R}_+^n} \psi_\infty^{q-1} u^2 dx'}{\int_{\partial \mathbb{R}_+^n} |u|^2 |x'|^{a-1} dx'}. \quad (16)$$

Substituting the explicit expression of $\psi_\infty(x) = V(\theta)r^{\frac{a-1}{q-1}}$ into (16), we have

$$J(q) = C_a - qV_B^{q-1}.$$

Definition 3.1 We say that an exponent q is JL-supercritical if $J(q) > 0$, JL-critical if $J(q) = 0$ and JL-subcritical if $J(q) < 0$.

Remark 2 Hence, an exponent q is JL-supercritical if $C_a > qV_B^{q-1}$, JL-critical if $C_a = qV_B^{q-1}$ and JL-subcritical if $C_a < qV_B^{q-1}$.

Next we will show that q is JL-supercritical if q and n are large enough, and q is JL-subcritical if q is close to $\frac{n-a}{n+a-2}$.

Lemma 3.1 There exists $n_0 \in \mathbb{N}$ and $q_1 > \frac{n_0-a}{n_0+a-2}$ such that for $n \geq n_0$ and $q > q_1$ we have $C_a > qV_B^{q-1}$.

Proof. Integrating (5) over $(0, \frac{\pi}{2})$, we have

$$V_B^q = \lim_{\theta \rightarrow \frac{\pi}{2}} V_\theta(\theta) \cos^a \theta = \gamma \int_0^{\frac{\pi}{2}} V(\theta) d\mu \leq \gamma \int_0^{\frac{\pi}{2}} V_B d\mu, \quad (17)$$

where in the last inequality we used the fact that $V_\theta > 0$. Hence we have

$$qV_B^{q-1} \leq q\gamma \int_0^{\frac{\pi}{2}} d\mu = qm_q(n+a-2-m_q)I_{n,a},$$

where $I_{n,a} := \int_0^{\frac{\pi}{2}} d\mu = \int_0^{\frac{\pi}{2}} \sin^{n-2} \cos^a \theta d\theta \rightarrow 0$ as $n \rightarrow +\infty$. Note that for $q > q_1$ we have $qm_q = q^{\frac{1-a}{q-1}} \leq C$, where C is a positive constant independence of n . On the other hand, by Lemma 2.2, one has $C_a > \frac{n+a-3}{2}$. So $qm_q(n+a-2-m_q)I_{n,a} < C_a$ for large n . Hence the result of this lemma is true.

Lemma 3.2 For $n \geq 3$, there exists $q_0 \geq \frac{n-a}{n+a-2}$ such that $C_a < qV_B^{q-1}$ for $q \in [\frac{n-a}{n+a-2}, q_0)$. Furthermore, in these low dimensions $n = 3, 4, 5, 6$, we have $C_a < qV_B^{q-1}$ for any $q \geq \frac{n-a}{n+a-2}$.

Proof. By the explicit expression of γ and $h_{n,a}$, it is easily seen that $h_{n,a} \geq \gamma$ for $q \in [\frac{n-a}{n+a-2}, +\infty)$, moreover, the strict inequality holds unless $q = \frac{n-a}{n+a-2}$. Let $e(\theta)$ be a positive solution of (13) with $e_B = V_B$. We set $W(\theta) := e(\theta) - V(\theta)$, then we have

$$\begin{cases} W_{\theta\theta} + (n-2) \cot \theta W_\theta - a \tan \theta W_\theta \geq \gamma W & \text{in } (0, \frac{\pi}{2}), \\ W = 0 & \text{on } \{\frac{\pi}{2}\}. \end{cases} \quad (18)$$

We claim that

$$W(\theta) \leq 0 \quad \text{in } (0, \frac{\pi}{2}). \quad (19)$$

First we suppose that $W(0) > 0$. Note that $\lim_{\theta \rightarrow 0} \frac{W_\theta(\theta)}{\tan \theta} = \lim_{\theta \rightarrow 0} \frac{W_\theta(\theta) - W_\theta(0)}{\theta} = W_{\theta\theta}(0)$. By this and (18), we have $(n-1)W_{\theta\theta}(0) \geq \gamma W(0) > 0$. Since $W_\theta(0) = 0$, there exists $\delta > 0$ such that $W_\theta > 0$ in $(0, \delta)$. Due to $W(\frac{\pi}{2}) = 0$, there exists $\theta_0 \in (0, \frac{\pi}{2})$ such that $W(\theta_0) > 0$, $W_\theta(\theta_0) = 0$ and $W_{\theta\theta}(\theta_0) \leq 0$. However, these contradict with (18). Hence we obtain $W(0) \leq 0$. Now we suppose (19) is false. Then there exists $\theta_1 \in (0, \frac{\pi}{2})$ such that $W(\theta_1) > 0$, $W_\theta(\theta_1) = 0$ and $W_{\theta\theta}(\theta_1) \leq 0$, which contradict with (18) again. Hence the claim (19) is proved.

Multiplying the equation in the following problem

$$-\Delta_{S,a} e = h_{n,a} e, \quad \theta \in (0, \frac{\pi}{2}), \quad \lim_{\theta \rightarrow \frac{\pi}{2}} e_\theta(\theta) \cos^a \theta = C_a e_B,$$

by $\sin^{n-2} \cos^a \theta$ and integrating over $(0, \frac{\pi}{2})$, we obtain

$$C_a e_B = h_{n,a} \int_0^{\frac{\pi}{2}} e(\theta) d\mu. \quad (20)$$

From (17), (20) and the assumption $e_B = V_B$, we have

$$qV_B^{q-1} - C_a = V_B^{-1}(qV_B^q - C_a e_B) = V_B^{-1} \left(q\gamma \int_0^{\frac{\pi}{2}} V(\theta) d\mu - h_{n,a} \int_0^{\frac{\pi}{2}} e(\theta) d\mu \right).$$

From (19) we have

$$qV_B^{q-1} - C_a \geq V_B^{-1}(q\gamma - h_{n,a}) \int_0^{\frac{\pi}{2}} e(\theta) d\mu.$$

Then to prove $qV_B^{q-1} > C_a$, we need to show $q\gamma > h_{n,a}$. From the explicit expression of γ , we have

$$q\gamma - h_{n,a} = \left(1 - a + \frac{1-a}{q-1}\right) \left((n+a-2) - \frac{1-a}{q-1} \right) - \left(\frac{n+a-2}{2} \right)^2. \quad (21)$$

Now we choose $q = \frac{n-a}{n+a-2}$ in (21), then we see that $q\gamma - h_{n,a} > 0$. Hence by a continuity principle, there exists $q_0 > \frac{n-a}{n+a-2}$ such that $q\gamma - h_{n,a} > 0$ for $q \in [\frac{n-a}{n+a-2}, q_0)$, which shows the first statement.

Next we prove the second statement. We regard the right-hand side of (21) as a function G of $\frac{1-a}{q-1}$. Then $q \geq \frac{n-a}{n+a-2}$ is equivalent to $\frac{1-a}{q-1} \in (0, \frac{n+a-2}{2}]$. Hence

$$\inf_{q \geq \frac{n-a}{n+a-2}} (q\gamma - h_{n,a}) = \inf_{\tau \in (0, \frac{n+a-2}{2}]} G(\tau), \quad (22)$$

where $G(\tau) := (1-a+\tau)((n+a-2)-\tau) - \frac{(n+a-2)^2}{4}$. Elementary computation shows that, for $n = 3$

$$\inf_{\tau \in (0, \frac{n+a-2}{2}]} G(\tau) = G(\frac{n+a-2}{2}) = \frac{1-a^2}{2} > 0,$$

and for $n = 4$

$$\inf_{\tau \in (0, \frac{n+a-2}{2}]} G(\tau) = G(\frac{n+a-2}{2}) = \frac{(2+a)(1-a)}{2} > 0.$$

For the case $n = 5$, if $a \in (-1, -\frac{1}{3}]$, we have

$$\inf_{\tau \in (0, \frac{n+a-2}{2}]} G(\tau) = G(\frac{n+a-2}{2}) = \frac{(3+a)(1-a)}{2} > 0,$$

and if $a \in (-\frac{1}{3}, 0)$, we have

$$\inf_{\tau \in (0, \frac{n+a-2}{2}]} G(\tau) = G(0) = \frac{(3+a)(1-a)}{4} > 0.$$

For the case $n = 6$, if $a \in (-1, -\frac{2}{3}]$, we have

$$\inf_{\tau \in (0, \frac{n+a-2}{2}]} G(\tau) = G(\frac{n+a-2}{2}) = \frac{(4+a)(1-a)}{2} > 0,$$

and if $a \in (-\frac{2}{3}, 0)$, we have

$$\inf_{\tau \in (0, \frac{n+a-2}{2}]} G(\tau) = G(0) = \frac{(4+a)(-a)}{4} > 0.$$

As for the case $n \geq 7$, we have $\inf_{\tau \in (0, \frac{n+a-2}{2}]} G(\tau) = G(0) < 0$.

The proof of this lemma is complete.

4 Estimate of eigenvalues

In this section we will make some estimates for the eigenvalues of the following problem.

$$-\Delta_{S,a}e = \lambda e, \quad \theta \in (0, \frac{\pi}{2}), \quad \lim_{\theta \rightarrow \frac{\pi}{2}} e_\theta(\theta) \cos^a \theta = qV_B^{q-1}e_B. \quad (23)$$

We denote by λ_i , $e_i(\theta)$ the i -th normalized eigenfunction such that

$$\int_0^{\frac{\pi}{2}} e_i(\theta) e_j(\theta) d\mu = \delta_{ij}, \quad e_{iB} > 0.$$

First we give estimates for the first eigenvalue for this problem.

Lemma 4.1 $\lambda_1 < -\gamma$ and

$$\begin{aligned} \lambda_1 &> -\frac{(n+a-2)^2}{4} && \text{if } q \text{ is } JL\text{-supercritical;} \\ \lambda_1 &= -\frac{(n+a-2)^2}{4} && \text{if } q \text{ is } JL\text{-critical;} \\ \lambda_1 &< -\frac{(n+a-2)^2}{4} && \text{if } q \text{ is } JL\text{-subcritical.} \end{aligned}$$

Proof. For the first eigenvalue λ_1 of (23), it is characterized by

$$\lambda_1 = \inf_{e \in \mathcal{H}_a^1(S_+^{n-1})} \frac{\|\partial_\theta e\|_{2,a}^2 - qV_B^{q-1}e_B^2}{\|e\|_{2,a}^2}. \quad (24)$$

Recall that $V(\theta)$ is a positive solution of (5), then $-\gamma$ is the first eigenvalue of the problem

$$-\Delta_{S,a}e = \lambda e, \quad \theta \in (0, \frac{\pi}{2}), \quad \lim_{\theta \rightarrow \frac{\pi}{2}} e_\theta(\theta) \cos^a \theta = V_B^{q-1}e_B.$$

Hence

$$-\gamma = \inf_{e \in \mathcal{H}_a^1(S_+^{n-1})} \frac{\|\partial_\theta e\|_{2,a}^2 - V_B^{q-1}e_B^2}{\|e\|_{2,a}^2}. \quad (25)$$

By (24), (25) and the fact $q > 1$, we obtain that $\lambda_1 < -\gamma$.

In Section 2 we have proved that $h_{n,a} = -\frac{(n+a-2)^2}{4}$ is the first eigenvalue of (11), hence we have

$$-\frac{(n+a-2)^2}{4} = \inf_{e \in \mathcal{H}_a^1(S_+^{n-1})} \frac{\|\partial_\theta e\|_{2,a}^2 - C_a e_B^2}{\|e\|_{2,a}^2}. \quad (26)$$

By Remark 2, (24) and (26), we obtain the rest results of this lemma.

We set

$$\sigma := (n+a-2) - 2m_q.$$

Elementary computation shows that $\sigma^2 + 4(\gamma + \lambda_1) = (n+a-2)^2 + 4\lambda_1$. From this and Lemma 4.1, we obtain the following lemma.

Lemma 4.2 *One has*

$$\begin{aligned}\sigma^2 + 4(\gamma + \lambda_1) &> 0 && \text{if } q \text{ is } JL\text{-supercritical;} \\ \sigma^2 + 4(\gamma + \lambda_1) &= 0 && \text{if } q \text{ is } JL\text{-critical;} \\ \sigma^2 + 4(\gamma + \lambda_1) &< 0 && \text{if } q \text{ is } JL\text{-subcritical.}\end{aligned}$$

Next we show that the second eigenvalue of (23) is positive.

Lemma 4.3 *It holds that $\lambda_2 > 0$.*

Proof. Let $e_2(\theta)$ be a corresponding eigenfunction with $e_2(0) < 0$. By Sturm's comparison theorem, $e_2(\theta)$ has just one zero in $(0, \frac{\pi}{2})$, which is denoted by θ_0 . Then $\partial_\theta e_2(\theta_0) \geq 0$. Clearly $\partial_\theta e_2(\theta_0) \neq 0$. Hence $\partial_\theta e_2(\theta_0) > 0$. Multiplying (23) by $\sin^{n-2} \theta_0 \cos^a \theta_0$ and integrating over $(0, \theta_0)$, we obtain

$$\partial_\theta e_2(\theta_0) = \frac{-\lambda_2}{\sin^{n-2} \theta_0 \cos^a \theta_0} \int_0^{\theta_0} e_2(\theta) d\mu,$$

which gives that $\lambda_2 > 0$.

5 Decay estimates and Limit behavior

In this section we will prove Theorem 1.1.

Recall that from Proposition 2.1, (2) admits a positive axial symmetric solution $u(x)$ satisfying $u(0) = 1$. We set $u_1(x) := u(x)$ and

$$u_\beta(x) := \beta u_1(\beta^{\frac{q-1}{1-a}} x).$$

Then $u_\beta(x)$ is a positive axial symmetric solution of (2) satisfying $u_\beta(0) = \beta$. Hence Theorem 1.1 (i) is proved.

Now we will follow the idea in [11] to prove Theorem 1.1 (ii). We denote a positive axial symmetric solution of (2) as $u(x) = u(r, \theta)$, which satisfies $u_\theta > 0$. Then we need to prove the decay estimate

$$u(r, \theta) = u(x) \leq C(1 + |x|)^{-m_q}. \quad (27)$$

We define

$$v(t, \theta) := r^{\frac{1-a}{q-1}} u(r, \theta), \quad r := e^t.$$

Then $v(t, \theta)$ satisfies

$$\begin{cases} v_{tt} + \sigma v_t - \gamma v + \Delta_{S,a} v = 0 & \text{in } \mathbb{R} \times (0, \frac{\pi}{2}), \\ \lim_{\theta \rightarrow \frac{\pi}{2}} v_\theta \cos^a \theta = v_B^q(t) & \text{on } \mathbb{R}, \end{cases} \quad (28)$$

where σ is the positive constant defined in the previous section. To prove the decay estimate (27), we need to show that $v(t, \theta)$ is bounded on $\mathbb{R}_+ \times (0, \frac{\pi}{2})$. We set

$$\tilde{v}(t) := \int_0^{\frac{\pi}{2}} v(t, \theta) d\mu.$$

We will first show that $\tilde{v}(t)$ is bounded on $\mathbb{R}_+$.

Multiplying (28) by $\sin^{n-2} \theta \cos^a \theta$ and integrating with respect to θ , we obtain

$$\tilde{v}_{tt} + \sigma \tilde{v}_t - \gamma \tilde{v} + v_B^q = 0. \quad (29)$$

For problem (29), the following lemma is obtained in [11].

Lemma 5.1 *Let $v(t, \theta)$ be a positive axial symmetric solution of (28) satisfying $v_\theta > 0$. Then $\tilde{v}(t)$ is bounded on $\mathbb{R}_+$.*

In order to prove that $v(t, \theta)$ is bounded on $\mathbb{R}_+ \times (0, \frac{\pi}{2})$, we need to establish an apriori estimate for the following problem

$$-\operatorname{div}(x_n^a \nabla u) + c(x) x_n^a u = 0 \quad \text{in } B_1^+, \quad \frac{\partial u}{\partial \nu^a} = F(x') u \quad \text{on } \partial D_1, \quad (30)$$

where $B_1^+ := \{x \in \mathbb{R}_+^n : |x| < 1\}$, $D_1 := \{x \in \partial \mathbb{R}_+^n : |x| < 1\}$. We will borrow the method used in Theorem 8.17 in [8] (see also in [11]) to prove the following apriori estimate.

Lemma 5.2 *Let $u(x) \in H_a^1(B_1^+)$ be a weak solution of (30) with $F \in L^p(D_1)$ for some $p > \frac{n-1}{1-a}$. Then there exists $C > 0$ depending on n , a , $\|F\|_{L^p(D_1)}$ and $\|c\|_{L^\infty(B_1^+)}$ such that*

$$\|u\|_{L^\infty(B_{1/2}^+)} \leq C \|u\|_{L_a^2(B_1^+)}.$$

Proof. Let $\chi(|x|)$ be a smooth cut-off function satisfying $\chi(|x|) = 0$ for $|x| \geq 1$ and set $\tilde{c} = \|c\|_{L^\infty(B_1^+)}$. We denote $u_+ = \max\{u, 0\}$, $u_- = \max\{-u, 0\}$. Multiplying the equation in (30) by $u_+^k \chi^2$ ($k > 1$) and integrating over B_1^+ , we obtain

$$\begin{aligned} & \frac{4k}{(k+1)^2} \int_{B_1^+} |\nabla u_+^{(k+1)/2}|^2 \chi^2 x_n^a dx \\ & \leq \int_{B_1^+} u_+^k |\nabla u_+| |\nabla \chi^2| x_n^a dx + \tilde{c} \int_{B_1^+} u_+^{k+1} \chi^2 x_n^a dx + \int_{D_1} F(x') u_+^{k+1} \chi^2 dx'. \end{aligned} \quad (31)$$

Let $p' = \frac{p}{p-1}$ be the dual exponent of p . Since $p > \frac{n-1}{1-a}$, then $2 < 2p' < \frac{2(n-1)}{n+a-2}$. We denote $2_* := \frac{2(n-1)}{n+a-2}$. Then there exists a $\varsigma \in (0, 1)$ such that $\frac{1}{2p'} = \frac{\varsigma}{2} + \frac{1-\varsigma}{2_*}$. By the Hölder inequality and an interpolation inequality, we have

$$\begin{aligned} & \int_{D_1} F(x') u_+^{k+1} \chi^2 dx' \leq \|F\|_{L^p(D_1)} \|u_+^{(k+1)/2} \chi\|_{L^{2p'}(D_1)}^2 \\ & \leq \|F\|_{L^p(D_1)} \left[\varepsilon \|u_+^{(k+1)/2} \chi\|_{L^{2_*}(D_1)} + \varepsilon^{-\frac{1-\varsigma}{\varsigma}} \|u_+^{(k+1)/2} \chi\|_{L^2(D_1)} \right]^2. \end{aligned}$$

Since $a \in (-1, 0)$, the following trace inequalities hold

$$\|w\|_{L^{2*}(D_1)} \leq C\|w\|_{H_a^1(B_1^+)}, \quad \|w\|_{L^2(D_1)} \leq A(\varepsilon, \varsigma)\|\nabla w\|_{L_a^2(B_1^+)} + \frac{C}{A(\varepsilon, \varsigma)}\|w\|_{L_a^2(B_1^+)},$$

where $A(\varepsilon, \varsigma) > 0$ is chosen small enough as $\varepsilon^{-\frac{1-\varsigma}{\varsigma}} A(\varepsilon, \varsigma) < \varepsilon$. Hence we obtain

$$\int_{D_1} F(x') u_+^{k+1} \chi^2 dx' \leq \|F\|_{L^p(D_1)} \left[4\varepsilon^2 \|u_+^{(k+1)/2} \chi\|_{H_a^1(B_1^+)} + C(\varepsilon) \|u_+^{(k+1)/2} \chi\|_{L_a^2(B_1^+)} \right]. \quad (32)$$

From (31) and (32), choosing $\varepsilon^2 = (\|F\|_{L^p(D_1)})^{-1} \frac{k}{2(k+1)^2}$, we obtain

$$\int_{B_1^+} |\nabla(u_+^{(k+1)/2} \chi)|^2 x_n^a dx \leq C(n, a, \|F\|_{L^p(D_1)}, \tilde{c})(k+1)^2 \int_{B_1^+} u_+^{k+1} (\chi + |\nabla \chi|)^2 x_n^a dx.$$

From [10], we know that there exists some $v > 2$ such that for $2^* := \frac{2v}{v-2}$, one has

$$\|u_+^{(k+1)/2} \chi\|_{L_a^{2^*}(B_1^+)} \leq C(n, a, \|F\|_{L^p(D_1)}, \tilde{c})(k+1) \|u_+^{(k+1)/2} (\chi + |\nabla \chi|)\|_{L_a^2(B_1^+)}. \quad (33)$$

It is now desirable to specify the cut-off function χ more precisely. Let r_1, r_2 be such that $\frac{1}{2} \leq r_1 \leq r_2 \leq \frac{3}{4}$ and set $\chi \equiv 1$ in $B_{r_1}^+$, $\chi \equiv 0$ in $B_1^+ \setminus B_{r_2}^+$ with $|\nabla \chi| \leq \frac{2}{r_2 - r_1}$. Writing $\eta := \frac{v}{v-2}$, we then have from (33)

$$\|u_+^{(k+1)/2}\|_{L_a^{2\eta}(B_{r_1}^+)} \leq C(k+1) \|u_+^{(k+1)/2}\|_{L_a^2(B_{r_2}^+)}. \quad (34)$$

Let us introduce

$$\Phi(\delta, r) := \left(\int_{B_r^+} |u_+|^\delta x_n^a dx \right)^{\frac{1}{\delta}}.$$

Due to the fact that $\int_{B_r^+} x_n^a dx < \infty$, it is verified that

$$\Phi(+\infty, r) = \lim_{\delta \rightarrow +\infty} \Phi(\delta, r) = \sup_{B_r^+} u_+, \quad \Phi(-\infty, r) = \lim_{\delta \rightarrow -\infty} \Phi(\delta, r) = \inf_{B_r^+} u_+.$$

From (34) we have

$$\Phi(\eta(k+1), r_1) \leq C(k+1)^{\frac{2}{(k+1)}} \Phi((k+1), r_2). \quad (35)$$

This inequality can now be iterated to yield the desired estimates. Hence, taking $\delta > 1$, we set $k+1 = \eta^i \delta$ and $r_i = 2^{-1} + 2^{-1-i}$, $i = 0, 1, \dots$, so that, by (35) we have

$$\Phi(\eta^i \delta, \frac{1}{2}) \leq (C\eta)^{2\sum_{i=0}^{i-1} \eta^{-i}} \Phi(\delta, 1) = \tilde{C} \Phi(\delta, 1),$$

where $\tilde{C}$ depending on $\|F\|_{L^p(D_1)}, \tilde{c}, a, n, \delta$. Consequently, letting i tend to infinity, we have

$$\|u_+\|_{L^\infty(B_{1/2}^+)} \leq C \|u_+\|_{L_a^\delta(B_1^+)}.$$

Let us now choose $\delta = 2$, then

$$\|u_+\|_{L^\infty(B_{1/2}^+)} \leq C \|u_+\|_{L_a^2(B_1^+)}.$$

By the same way, we can obtain the estimate for u_- .

We complete the proof of this lemma.

Remark 3 *Similar estimate as that of Lemma 5.2 had been obtained by Fabes-Kenig-Serapioni [7] (Corollary 2.3.4).*

Lemma 5.3 *Let $v(t, \theta)$ be a positive axial symmetric solution of (28) satisfying $v_\theta > 0$. Then $v(t, \theta)$ is bounded on $\mathbb{R}_+ \times (0, \frac{\pi}{2})$.*

Proof. We apply a test function method to prove a boundedness of $v(t, \theta)$. Let $\chi(t)$ be a cut-off function with a compact support in $\mathbb{R}_+$. Multiplying (28) by $v^{-\frac{1}{2}}(t, \theta) \chi^2 \sin^{n-2} \theta \cos^a \theta$ and integrating on $\mathbb{R}_+ \times (0, \frac{\pi}{2})$, we obtain

$$\int_{\mathbb{R}} \left(\int_0^{\frac{\pi}{2}} v^{-\frac{3}{2}} (v_t^2 + v_\theta^2) d\mu + v_B^{q-1/2} \right) \chi^2 dt \leq C \int_{\mathbb{R}} \int_0^{\frac{\pi}{2}} v^{\frac{1}{2}} (\chi^2 + \chi_t^2) d\mu dt. \quad (36)$$

We need to use the following trace inequality for a two dimensional domain $(a_1, a_2) \times (b_1, b_2)$

$$\int_{a_1}^{a_2} |\phi(\xi_1, b_2)|^\alpha d\xi_1 \leq C_\alpha \left(\int_{a_1}^{a_2} \int_{b_1}^{b_2} [\phi^2 + \phi_{\xi_1}^2 + \phi_{\xi_2}^2] d\xi_2 d\xi_1 \right)^{\frac{\alpha}{2}},$$

where $\alpha \geq 1$ and $C_\alpha > 0$ is a constant depending on α , $a_2 - a_1$ and $b_2 - b_1$. Applying this inequality with $\phi(\xi_1, \xi_2) = v^{\frac{1}{4}}(\xi_1, \xi_2)$, $a_1 = \tau - 1$, $a_2 = \tau + 1$, $b_1 = \frac{\pi}{4}$, $b_2 = \frac{\pi}{2}$, then we obtain

$$\int_{\tau-1}^{\tau+1} |v_B^{\frac{1}{4}}|^\alpha dt \leq C_\alpha \left(\int_{\tau-1}^{\tau+1} \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} [v^{\frac{1}{2}} + v^{-\frac{3}{2}} (v_t^2 + v_\theta^2)] d\theta dt \right)^{\frac{\alpha}{2}}.$$

We take a cut-off function $\chi(t)$ such that $\chi(t) = 1$ if $t \in [\tau - 1, \tau + 1]$ and $\chi(t) = 0$ if $t \in \mathbb{R} \setminus (\tau - 2, \tau + 2)$. It is clear that $\sin \theta \geq \frac{\sqrt{2}}{2}$ for $\theta \in [\frac{\pi}{4}, \frac{\pi}{2}]$, and $\cos^a \theta > 1$, since $a \in (-1, 0)$.

Hence from (36), for $\alpha \geq 1$, we have

$$\int_{\tau-1}^{\tau+1} |v_B^{\frac{1}{4}}|^\alpha dt \leq C_\alpha \left(\int_{\tau-2}^{\tau+2} \int_0^{\frac{\pi}{2}} v^{\frac{1}{2}} d\mu dt \right)^{\frac{\alpha}{2}} \leq C_\alpha \left(\int_{\tau-2}^{\tau+2} \tilde{v} dt \right)^{\frac{\alpha}{4}},$$

where in the last inequality we have used Hölder inequality. Therefore, from Lemma 5.1, for $\alpha \geq 1$ there exists $\bar{C}_\alpha > 0$ independent of $\tau > 0$ such that

$$\int_{\tau-1}^{\tau+1} |v_B^{\frac{1}{4}}|^\alpha dt \leq \bar{C}_\alpha.$$

Then from Lemma 5.2, we deduce that $v(t, \theta)$ is bounded on $\mathbb{R}_+ \times (0, \frac{\pi}{2})$.

The proof of this lemma is complete.

In the rest of this section, we will prove Theorem 1.1 (iii): the limit behavior of positive axial symmetric solutions.

Recall that $v(t, \theta)$ satisfies (28). Note that $\sigma, \gamma > 0$ if $q > \frac{n-a}{n+a-2}$. We define the following energy function $E(t)$ associated with (28) by

$$E(t) := \frac{1}{2} \|\partial_t v\|_{2,a}^2 - \frac{\gamma}{2} \|v\|_{2,a}^2 - \frac{1}{2} \|\partial_\theta v\|_{2,a}^2 + \frac{1}{q+1} (v_B(t))^{q+1}.$$

Simple calculation shows that

$$\partial_t E(t) = -\sigma \|\partial_t v\|_{2,a}^2.$$

Lemma 5.4 *Let $q > \frac{n-a}{n+a-2}$ and $v(t, \theta)$ be a positive bounded solution of (28) with $\lim_{t \rightarrow -\infty} E(t) = 0$. Then one has*

$$\lim_{t \rightarrow +\infty} v(t, \theta) = V(\theta) \quad \text{in } C[0, \pi/2].$$

Proof. From the boundary condition in (28), $a \in (-1, 0)$ and the result that $v(t, \theta)$ is bounded on $\mathbb{R} \times (0, \pi/2)$, we have $v_\theta(t, \frac{\pi}{2}) = 0$. The elliptic regularity theory assures a boundedness of $\partial_t v(t, \theta)$ and $\partial_\theta v(t, \theta)$. Hence we deduce that $E(t)$ is bounded and

$$0 < \sigma \int_{-\infty}^{\infty} \int_0^{\frac{\pi}{2}} v_t^2 d\mu dt = \lim_{t \rightarrow -\infty} E(t) - \lim_{t \rightarrow \infty} E(t) = - \lim_{t \rightarrow \infty} E(t) < \infty. \quad (37)$$

Let $\{t_i\}_{i \in \mathbb{N}}$ be any sequence satisfying $\lim_{i \rightarrow \infty} t_i = +\infty$ and set $v_i(t, \theta) = v(t + t_i, \theta)$. Then there exists a subsequence of $\{t_i\}_{i \in \mathbb{N}}$, which is still denoted by the same symbol such that $v_i(t, \theta)$ converges to some function $v_\infty(t, \theta)$ in $C([-1, 1] \times [0, \pi/2])$. By (37) we find that $\partial_t v_\infty(t, \theta) \equiv 0$. Then we denote $v_\infty(t, \theta) =: v_\infty(\theta)$ and $v_\infty(\theta)$ satisfies

$$-\gamma v + \Delta_{S,a} v = 0 \quad \text{in } (0, \pi/2), \quad \lim_{\theta \rightarrow \frac{\pi}{2}} v_\theta \cos^a \theta = v_B^q.$$

By $\lim_{t \rightarrow -\infty} E(t) = 0$ and (37), we have $\lim_{t \rightarrow \infty} E(t) < 0$, which yields that the limiting function $v_\infty(\theta)$ is not a trivial one. Hence by Proposition 2.2, we deduce that $v_\infty(\theta) \equiv V(\theta)$. Therefore we obtain the result that $\lim_{i \rightarrow \infty} v(t_i, \theta) = V(\theta)$ in $C[0, \pi/2]$ for any sequence $\{t_i\}_{i \in \mathbb{N}}$ converging to $+\infty$. Hence

$$\lim_{t \rightarrow \infty} v(t, \theta) = V(\theta) \quad \text{in } C[0, \pi/2].$$

The lemma is proved.

Proof of Theorem 1.1 (iii) Recalling that $\psi_\infty(x) = V(\theta) r^{\frac{a-1}{q-1}}$ and $u_\beta(x) = \beta u_1(\beta^{\frac{q-1}{1-a}} x)$, we have

$$|u_\beta(x) - \psi_\infty(x)| = \left| \beta u_1(\beta^{\frac{q-1}{1-a}} x) - V(\theta) |x|^{\frac{a-1}{q-1}} \right| = \left| v_1(\beta^{\frac{q-1}{1-a}} x) - V(\theta) \right| |x|^{\frac{a-1}{q-1}}.$$

Note that $\lim_{t \rightarrow -\infty} v(t, \theta) = 0$, since u is bounded. Then the condition $\lim_{t \rightarrow -\infty} E(t) = 0$ in Lemma 5.4 is true. Hence by Lemma 5.4, we have

$$\lim_{\beta \rightarrow +\infty} |u_\beta(x) - \psi_\infty(x)| = 0, \quad x \neq 0.$$

We complete the proof of Theorem 1.1 (iii).

6 Asymptotic expansion

In this section, we will prove Theorem 1.2. We need to study the asymptotic behavior of

$$w(t, \theta) = V(\theta) - v(t, \theta).$$

Note that $w(t, \theta)$ satisfies

$$\begin{cases} w_{tt} + \sigma w_t - \gamma w + \Delta_{S,a} w = 0 & \text{in } \mathbb{R} \times (0, \frac{\pi}{2}), \\ \lim_{\theta \rightarrow \frac{\pi}{2}} w_\theta \cos^a \theta = q v_B^{q-1} w_B + g(w_B) & \text{on } \mathbb{R}, \end{cases} \quad (38)$$

where $g(w)$ is given by

$$g(w) = v_B^q - (v_B - w)^q - q v_B^{q-1} w.$$

Since $L_{\text{sym}}^2(S_+^{n-1})$ is spanned by the eigenfunctions $\{e_i(\theta)\}_{i \in N}$ of (23), then $w(t, \theta)$ can be expanded as

$$w(t, \theta) = \sum_{i=1}^{\infty} z_i(t) e_i(\theta), \quad t \in \mathbb{R}.$$

Multiplying the equation in (23) by $e_i(\theta) \sin^{n-2} \theta \cos^a \theta$ and integrating with respect to θ on $(0, \frac{\pi}{2})$, we obtain

$$z_i'' + \sigma z_i' - (\gamma + \lambda_i) z_i = f_i(t), \quad (39)$$

where $f_i(t) := -g(w_B(t)) e_{iB}$.

We first consider the case $i \geq 2$. Recall that Lemma 4.3 shows that $\lambda_i > 0$ for $i \geq 2$. Hence the corresponding quadratic equation

$$\rho^2 + \sigma \rho - (\gamma + \lambda_i) = 0 \quad (40)$$

to (39) admits two real roots

$$\rho_i^\pm = \frac{-\sigma \pm \sqrt{\sigma^2 + 4(\gamma + \lambda_i)}}{2}. \quad (41)$$

Note that $\rho_i^- < 0 < \rho_i^+$. From [11] we know that for $i \geq 2$

$$\begin{aligned} z_i(t) &= z_i(0) e^{\rho_i^- t} - \frac{e^{\rho_i^- t}}{\sqrt{\sigma^2 + 4(\gamma + \lambda_i)}} \int_0^t (e^{-\rho_i^- s} - e^{-\rho_i^+ s}) f_i(s) ds \\ &\quad - \frac{e^{\rho_i^+ t} - e^{\rho_i^- t}}{\sqrt{\sigma^2 + 4(\gamma + \lambda_i)}} \int_t^\infty e^{-\rho_i^+ s} f_i(s) ds. \end{aligned} \quad (42)$$

For the case $i = 1$, by Lemma 4.2, (39) admits two real roots $\rho_1^- < \rho_1^+ < 0$ if q is JL-supercritical, just one root $\rho_1 = -\frac{\sigma}{2}$ if q is JL-critical and admits no real roots if q is JL-subcritical. Hence we obtain for the JL-supercritical case

$$\begin{aligned} z_1(t) = & z_1(0)e^{\rho_1^- t} - \frac{z_1'(0) - \rho_1^- z_1(0)}{\sqrt{\sigma^2 + 4(\gamma + \lambda_1)}}(e^{\rho_1^+ t} - e^{\rho_1^- t}) \\ & + \int_0^t \frac{1 - e^{(2\rho_1^- + \sigma)(t-s)}}{\sqrt{\sigma^2 + 4(\gamma + \lambda_1)}} e^{\rho_1^+(t-s)} f_1(s) ds, \end{aligned} \quad (43)$$

for the JL-critical case

$$z_1(t) = z_1(0)e^{\rho_1 t} + (z_1'(0) - \rho_1 z_1(0))te^{\rho_1 t} + \int_0^t (t-s)e^{\rho_1(t-s)} f_1(s) ds \quad (44)$$

and for the JL-subcritical case

$$\begin{aligned} z_1(t) = & \frac{1}{K} \left(\frac{\sigma}{2} z_1(0) + z_1'(0) \right) (\sin Kt) e^{-\frac{\sigma}{2} t} + z_1(0) (\cos Kt) e^{-\frac{\sigma}{2} t} \\ & + \frac{1}{K} \int_0^t [(\sin Kt)(\cos Ks) - (\sin Ks)(\cos Kt)] e^{-\frac{\sigma(t-s)}{2}} f_1(s) ds, \end{aligned} \quad (45)$$

where K , given in Theorem 1.2, is the imaginary part of a root of $\rho^2 + \sigma\rho - (\gamma + \lambda_1) = 0$.

To prove Theorem 1.2, we need to establish the following proposition.

Proposition 6.1 (i) *If q is JL-supercritical, then there exists $\xi_1 > 0$ and $\varepsilon, C > 0$ such that*

$$\|w(t, \theta) - \xi_1 e^{\rho_1^+ t} e_1\|_\infty \leq C e^{(\rho_1^+ - \varepsilon)t}, \quad t > 0;$$

(ii) *If q is JL-critical, then there exists $\xi_1 > 0$, $\xi_2 \in \mathbb{R}$ and $\varepsilon, C > 0$ such that*

$$\|w(t, \theta) - (\xi_1 t + \xi_2) e^{-\sigma t/2} e_1\|_\infty \leq C e^{-(\sigma + \varepsilon)t/2}, \quad t > 0;$$

(iii) *If q is JL-subcritical, then one of the following two expansions holds.*

(iii-1) *there exist $(\xi_1, \xi_2) \neq 0$ and $\varepsilon, C > 0$ such that*

$$\|w(t, \theta) - (\xi_1 (\sin Kt) + \xi_2 (\cos Kt)) e^{-\sigma t/2} e_1\|_\infty \leq C e^{-(\sigma + \varepsilon)t/2}, \quad t > 0;$$

(iii-2) *there exist $\xi \neq 0$ and $\varepsilon, C > 0$ such that*

$$\|w(t, \theta) - \xi e^{\rho_2^- t} e_2\|_\infty \leq C e^{(\rho_2^- - \varepsilon)t}, \quad t > 0.$$

As a consequence of Proposition 6.1, we immediately obtain Theorem 1.2.

Proof of Theorem 1.2. We only show the proof of JL-supercritical case in Theorem 1.2, since the proof of the rest part is similar. Let q be JL-supercritical and set $\rho_1 = |\rho_1^+|$. By Proposition 6.1 we obtain

$$\|v(t, \theta) - (V - \xi_1 e^{-\rho_1 t} e_1)\|_\infty \leq C e^{-(\rho_1 + \varepsilon)t}, \quad t > 0.$$

Going back to the original variable, we obtain

$$u(r, \theta) = V(\theta)r^{-m_q} - \xi_1 r^{-m_q - \rho_1} e_1(\theta) + O(r^{-m_q - \rho_1 - \varepsilon}), \quad r \gg 1.$$

Plugging the explicit expression of σ and γ into (41), we obtain the explicit expression of ρ_1 , which coincides with the expression in Theorem 1.2.

Now the remaining task for us is to prove Proposition 6.1.

First we show that $w(t, \theta)$ decays exponentially as $t \rightarrow +\infty$.

Lemma 6.1 *There exists $\varepsilon, C > 0$ such that*

$$\|w(t, \cdot)\|_{2,a} \leq C e^{-\varepsilon t}, \quad t > 0.$$

Proof. From (42), we obtain for $i \geq 2$

$$\begin{aligned} |z_i(t)| &\leq |z_i(0)|e^{\rho_i^- t} + \frac{e^{\rho_i^- t}}{\sqrt{\lambda_i}} \int_0^t e^{-\rho_i^- s} |f_i(s)| ds + \frac{e^{\rho_i^+ t}}{\sqrt{\lambda_i}} \int_t^\infty e^{-\rho_i^+ s} |f_i(s)| ds \\ &\leq |z_i(0)|e^{\rho_i^- t} + \frac{e^{\rho_i^- t/2}}{\sqrt{\lambda_i |\rho_i^-|}} \left(\int_0^t e^{-\rho_i^- s} |f_i(s)|^2 ds \right)^{1/2} \\ &\quad + \frac{e^{\rho_i^+ t/2}}{\sqrt{\lambda_i |\rho_i^+|}} \left(\int_t^\infty e^{-\rho_i^+ s} |f_i(s)|^2 ds \right)^{1/2}. \end{aligned}$$

By a trace inequality

$$|e_{iB}| \leq C \|e_i\|_{2,a}^{1/2} \|\partial_\theta e_i\|_{2,a}^{1/2},$$

we have

$$\begin{aligned} \|\partial_\theta e_i\|_{2,a}^2 &= 2\|\partial_\theta e_i\|_{2,a}^2 - \|\partial_\theta e_i\|_{2,a}^2 = 2(qV_B^{q-1} e_{iB}^2 + \lambda_i \|e_i\|_{2,a}^2) - \|\partial_\theta e_i\|_{2,a}^2 \\ &\leq 2\lambda_i \|e_i\|_{2,a}^2 + CqV_B^{q-1} \|e_i\|_{2,a} \|\partial_\theta e_i\|_{2,a} - \|\partial_\theta e_i\|_{2,a}^2 \\ &\leq 2\lambda_i \|e_i\|_{2,a}^2 + C\|e_i\|_{2,a}^2 \leq C\lambda_i \|e_i\|_{2,a}^2. \end{aligned}$$

Therefore, since $\|e_i\|_{2,a} = 1$, we deduce that $\|\partial_\theta e_i\|_{2,a}^2 \leq C\lambda_i$. Then we have

$$|e_{iB}| \leq C \|e_i\|_{2,a}^{1/2} \|\partial_\theta e_i\|_{2,a}^{1/2} \leq C\lambda_i^{\frac{1}{4}}.$$

Recalling that $f_i(t) = -g(w_B(t))e_{iB}$, we obtain

$$|f_i(s)| \leq C\lambda_i^{\frac{1}{4}} w_B^2(s) \leq C\lambda_i^{\frac{1}{4}} \|w(s, \cdot)\|_{2,a} \|\partial_\theta w(s, \cdot)\|_{2,a}.$$

Since $|\rho_i^-|, |\rho_i^+| \geq C\sqrt{\lambda_i}$ for some $C > 0$, we have for $i \geq 2$

$$\begin{aligned} |z_i(t)|^2 &\leq C|z_i(0)|^2 e^{2\rho_i^- t} + \frac{C}{\lambda_i} \left(\int_0^t e^{\rho_i^- (t-s)} \varphi(s) \|w(s, \cdot)\|_{2,a}^2 ds \right. \\ &\quad \left. + \int_t^\infty e^{-\rho_i^+ (s-t)} \varphi(s) \|w(s, \cdot)\|_{2,a}^2 ds \right) \\ &\leq C|z_i(0)|^2 e^{2\rho_i^- t} + \frac{C}{\lambda_i} \left(\int_0^t e^{\rho_i^- (t-s)} \varphi(s) \|w(s, \cdot)\|_{2,a}^2 ds \right. \\ &\quad \left. + \int_t^\infty e^{-\rho_i^+ (s-t)} \varphi(s) \|w(s, \cdot)\|_{2,a}^2 ds \right), \end{aligned}$$

where $\varphi(s) := \|\partial_\theta w(s, \cdot)\|_{2,a}^2$. Applying Strum's comparison theorem, we can obtain the following estimate of the eigenvalues of problem (23)

$$C_1 i^2 \leq \lambda_i \leq C_2 i^2, \quad \forall i > 1,$$

where C_1, C_2 are positive constants independence of i . Hence we have

$$\begin{aligned} \sum_{i=2}^{\infty} |z_i(t)|^2 &\leq C \|w(0, \cdot)\|_{2,a}^2 e^{2\rho_2^- t} + C \left(\int_0^t e^{\rho_2^- (t-s)} \varphi(s) \|w(s, \cdot)\|_{2,a}^2 ds \right. \\ &\quad \left. + \int_t^\infty e^{-\rho_2^+ (s-t)} \varphi(s) \|w(s, \cdot)\|_{2,a}^2 ds \right). \end{aligned}$$

Similarly, from (43)-(45), we obtain

$$|z_1(t)|^2 \leq \begin{cases} C e^{2\rho_1^+ t} + C \int_0^t e^{\rho_1^+ (t-s)} \varphi(s) \|w(s, \cdot)\|_{2,a}^2 ds & \text{if } q \text{ is JL-supercritical,} \\ C t^2 e^{2\rho_1 t} + C \int_0^t (t-s)^2 e^{\rho_1 (t-s)} \varphi(s) \|w(s, \cdot)\|_{2,a}^2 ds & \text{if } q \text{ is JL-critical,} \\ C e^{-\sigma t} + C \int_0^t e^{-\sigma (t-s)} \varphi(s) \|w(s, \cdot)\|_{2,a}^2 ds & \text{if } q \text{ is JL-subcritical.} \end{cases}$$

Hence there exists $\delta > 0$ such that

$$\|w(t, \cdot)\|_{2,a}^2 \leq C e^{-\delta t} + C \left(\int_0^t e^{-\delta (t-s)} \varphi(s) \|w(s, \cdot)\|_{2,a}^2 ds + \int_t^\infty e^{-\delta (s-t)} \varphi(s) \|w(s, \cdot)\|_{2,a}^2 ds \right)$$

By Lemma 6.2 stated below, we deduce the desired conclusion.

Lemma 6.2 ([11]) *Assume $\eta(t)$ and $\varphi(t)$ are positive continuous functions defined on $\overline{\mathbb{R}}_+$ converging to zero as $t \rightarrow +\infty$. Moreover $\eta(t)$ satisfies*

$$\eta(t) \leq C e^{-\delta t} + C \left(\int_0^t e^{-\delta (t-s)} \varphi(s) \eta(s) ds + \int_t^\infty e^{-\delta (s-t)} \varphi(s) \eta(s) ds \right), \quad t > 0$$

for some $\delta > 0$. Then there exists $\varepsilon > 0$ such that $\eta(t) \leq C e^{-\varepsilon t}$.

Next we will show more precise decay rates of $\|w(t, \cdot)\|_{2,a}$.

Lemma 6.3 *There exists $C > 0$ such that for $t > 0$*

$$\|w(t, \cdot)\|_{2,a} \leq \begin{cases} C e^{\rho_1^+ t} & \text{if } q \text{ is JL-supercritical,} \\ C(1+t) e^{\rho_1 t} & \text{if } q \text{ is JL-critical,} \\ C e^{-\sigma t/2} & \text{if } q \text{ is JL-subcritical.} \end{cases}$$

Proof. Repeating the argument given in the proof of Lemma 6.1, we have

$$\begin{aligned} \sum_{i=k}^{\infty} |z_i(t)|^2 &\leq C e^{2\rho_k^- t} + C \left(\int_0^t e^{\rho_k^- (t-s)} \varphi(s) \|w(s, \cdot)\|_{2,a}^2 ds \right. \\ &\quad \left. + \int_t^\infty e^{-\rho_k^+ (s-t)} \varphi(s) \|w(s, \cdot)\|_{2,a}^2 ds \right). \end{aligned}$$

Since $w(t, \cdot)$ is uniformly bounded in $\mathcal{H}_a^2(S_+^{n-1})$, by an interpolation inequality, it holds that

$$\|\partial_\theta w(t, \cdot)\|_{2,a}^2 \leq C \|w(t, \cdot)\|_{2,a} \|w(t, \cdot)\|_{\mathcal{H}_a^2(S_+^{n-1})} \leq C \|w(t, \cdot)\|_{2,a}.$$

Recalling $\varphi(s) = \|\partial_\theta w(s, \cdot)\|_{2,a}^2$, by Lemma 6.1, we have for $k \geq 2$

$$\sum_{i=k}^{\infty} |z_i(t)|^2 \leq C [e^{2\rho_k^- t} + e^{\rho_k^- t} + e^{-3\varepsilon t}].$$

From (42)-(45), we have that for $2 \leq i \leq k-1$

$$\begin{aligned} |z_i(t)| &\leq C e^{\rho_i^- t} + C e^{\rho_i^- t} \int_0^t e^{-\rho_i^- s} e^{-3\varepsilon s/2} ds + C e^{\rho_i^+ t} \int_t^\infty e^{-\rho_i^+ s} e^{-3\varepsilon s/2} ds \\ &\leq C [e^{\rho_i^- t} + e^{-3\varepsilon t/2}] \leq C [e^{\rho_2^- t} + e^{-3\varepsilon t/2}] \end{aligned}$$

and

$$|z_1(t)| \leq \begin{cases} C [e^{\rho_1^+ t} + e^{-3\varepsilon t/2}] & \text{if } q \text{ is JL-supercritical,} \\ C [(1+t)e^{\rho_1^+ t} + e^{-3\varepsilon t/2}] & \text{if } q \text{ is JL-critical,} \\ C [e^{-\sigma t/2} + e^{-3\varepsilon t/2}] & \text{if } q \text{ is JL-subcritical.} \end{cases} \quad (46)$$

We choose $k \geq 2$ large enough such that $|\rho_k^-| \geq 2|\rho_2^-|$, then we obtain

$$\sum_{i=2}^{\infty} |z_i(t)|^2 \leq C [e^{2\rho_2^- t} + e^{-3\varepsilon t}]. \quad (47)$$

We first consider the JL-supercritical case. If $\varepsilon \geq \frac{2|\rho_1^+|}{3}$, from (46)-(47) we know that this lemma is true. If $\varepsilon < \frac{2|\rho_1^+|}{3}$, from (46)-(47) we know that

$$\|w(t, \cdot)\|_{2,a} \leq C e^{-3\varepsilon t/2}.$$

Repeating the above procedure, if $\frac{2|\rho_1^+|}{3} > \varepsilon \geq \frac{4|\rho_1^+|}{9}$, from (46)-(47) (now the quantity ε in these inequalities should be changed into the quantity $\frac{3\varepsilon}{2}$), we know that this lemma is also true. If $\varepsilon < \frac{4|\rho_1^+|}{9}$, we know that

$$\|w(t, \cdot)\|_{2,a} \leq C e^{-9\varepsilon t/4}.$$

After repeating such procedure finite times, we can obtain

$$\|w(t, \cdot)\|_{2,a} \leq C e^{-\rho_1^+ t}.$$

Therefore we complete the proof of the JL-supercritical case. The JL-critical case and JL-subcritical case can be proved similarly.

Furthermore we see that the decay rate of $\|w(t, \cdot)\|_{2,a}$ given in Lemma 6.3 is exact one.

Lemma 6.4 *There exists $\xi_1, \xi_2 \in \mathbb{R}$ and $\varepsilon, C > 0$ such that for $t > 0$*

$$\begin{aligned} \|e^{-\rho_1^+ t} w(t, \cdot) - \xi_1 e_1\|_{2,a} &\leq C e^{-\varepsilon t} && \text{if } q \text{ is JL-supercritical,} \\ \|e^{\sigma t/2} w(t, \cdot) - (\xi_1 t + \xi_2) e_1\|_{2,a} &\leq C e^{-\varepsilon t} && \text{if } q \text{ is JL-critical,} \\ \|e^{\sigma t/2} w(t, \cdot) - (\xi_1 \sin(Kt) + \xi_2 \cos(Kt)) e_1\|_{2,a} &\leq C e^{-\varepsilon t} && \text{if } q \text{ is JL-subcritical.} \end{aligned} \quad (48)$$

Moreover the norm $\|\cdot\|_{2,a}$ can be replaced by the norm $\|\cdot\|_\infty$.

Proof. By Lemma 6.3 and (47), for the JL-supercritical case, we have

$$\sum_{i=2}^{\infty} |z_i(t)|^2 \leq C[e^{2\rho_2^- t} + e^{3\rho_1^+ t}].$$

Hence it is sufficient to estimate $z_1(t)$. Similarly, we also only need to estimate $z_1(t)$ in the JL-critical and JL-subcritical case.

We first consider the JL-supercritical case. We set $z_1(t) = e^{\rho_1^+ t} y_1(t)$. From (39), $y_1(t)$ solves

$$y_1'' + (2\rho_1^+ + \sigma)y_1' = e^{-\rho_1^+ t} f_1. \quad (49)$$

Then since $e^{(2\rho_1^+ + \sigma)t}[y_1'' + (2\rho_1^+ + \sigma)y_1'] = (e^{(2\rho_1^+ + \sigma)t} y_1')' = e^{(\rho_1^+ + \sigma)t} f_1$, we have

$$y_1'(t) = y_1'(0)e^{-(2\rho_1^+ + \sigma)t} + e^{-(2\rho_1^+ + \sigma)t} \int_0^t e^{(\rho_1^+ + \sigma)s} f_1(s) ds.$$

Hence by Lemma 6.3, there exists $\varepsilon > 0$ such that $|y_1'(t)| \leq C e^{-\varepsilon t}$, which yields that $\lim_{t \rightarrow +\infty} y_1(t)$ exists. We denote $\xi_1 := \lim_{t \rightarrow +\infty} y_1(t)$. So we have

$$|y_1(t) - \xi_1| \leq \int_t^\infty |y_1'(s)| ds \leq C e^{-\varepsilon t},$$

which shows (48) for the JL-supercritical case.

For the JL-critical case, we set $z_1(t) = t e^{\rho_1^+ t} y_1(t)$. From (39), $y_1(t)$ satisfies

$$t y_1'' + 2 y_1' = e^{-\rho_1^+ t} f_1. \quad (50)$$

This and the equation $t^2 y_1'' + 2t y_1' = (t^2 y_1')'$ give that

$$\begin{aligned} y_1'(t) &= \frac{1}{t^2} \int_0^t s e^{-\rho_1^+ s} f_1(s) ds \\ &= \frac{1}{t^2} \int_0^\infty s e^{-\rho_1^+ s} f_1(s) ds - \frac{1}{t^2} \int_t^\infty s e^{-\rho_1^+ s} f_1(s) ds, \end{aligned} \quad (51)$$

which gives that $y_1' \in L^1([1, \infty))$. So $\lim_{t \rightarrow +\infty} y_1(t)$ exists and we denote $\xi_1 := \lim_{t \rightarrow +\infty} y_1(t)$. We set $\xi_2 := \int_0^\infty s e^{-\rho_1^+ s} f_1(s) ds$. Then, from (51), we have

$$\left| y_1(t) - \xi_1 + \frac{\xi_2}{t} \right| \leq C e^{-\varepsilon t}, \quad t > 0$$

for some $\varepsilon > 0$, which shows (48) for the JL-critical case.

Finally we consider the JL-subcritical case. From (45), we have

$$\begin{aligned} z_1(t) &= \frac{1}{K} \left(\frac{\sigma}{2} z_1(0) + z_1'(0) \right) (\sin Kt) e^{-\frac{\sigma}{2}t} + z_1(0) (\cos Kt) e^{-\frac{\sigma}{2}t} \\ &\quad + \frac{e^{-\sigma t/2}}{K} \left((\sin Kt) \int_0^t (\cos Ks) e^{\sigma s/2} f_1(s) ds - (\cos Kt) \int_0^t (\sin Ks) e^{\sigma s/2} f_1(s) ds \right). \end{aligned}$$

We set $\alpha_1 := \int_0^t (\cos Ks) e^{\sigma s/2} f_1(s) ds$ and $\alpha_2 := \int_0^t (\sin Ks) e^{\sigma s/2} f_1(s) ds$. From Lemma 6.3, we deduce that $|\alpha_1|, |\alpha_2| < +\infty$. Therefore we obtain

$$\begin{aligned} &\left| z_1(t) - \frac{1}{K} \left(\frac{\sigma}{2} z_1(0) + z_1'(0) + \alpha_1 \right) (\sin Kt) e^{-\frac{\sigma}{2}t} - \left(z_1(0) - \frac{\alpha_2}{K} \right) (\cos Kt) e^{-\frac{\sigma}{2}t} \right| \\ &\leq \frac{1}{K} e^{-\frac{\sigma}{2}t} \int_t^\infty e^{\frac{\sigma}{2}s} |f_1(s)| ds, \end{aligned} \quad (52)$$

which shows (48) for the JL-subcritical case.

Now we consider the L^∞ -estimate. For the JL-supercritical case, we set $Y(t, \theta) := e^{-\rho_1^+ t} w(t, \theta) - \xi_1 e_1(\theta)$. Then we have

$$\begin{cases} Y_{tt} + (2\rho_1^+ + \sigma)Y_t + \lambda_1 Y + \Delta_{S,a} Y = 0 & \text{in } \mathbb{R} \times (0, \frac{\pi}{2}), \\ \lim_{\theta \rightarrow \frac{\pi}{2}} Y_\theta \cos^a \theta = qv_B^{q-1} Y_B + e^{-\rho_1^+ t} g(w_B) & \text{on } \mathbb{R}. \end{cases} \quad (53)$$

By the similar argument as given in Lemma 5.2, we know that there exists $C > 0$ such that

$$\sup_{t-1 \leq \tau \leq t+1} \|Y(\tau, \cdot)\|_\infty \leq C \left(\int_{t-2}^{t+2} \|Y(\tau, \cdot)\|_{2,a}^2 d\tau \right)^{\frac{1}{2}} + C \sup_{t-2 \leq \tau \leq t+2} e^{-\rho_1^+ \tau} |g(w_B(\tau))|.$$

Since $|g(w_B(\tau))| \leq C w_B^2(\tau)$, by a trace inequality and an interpolation, we have

$$|g(w_B(t))| \leq C w_B^2(t) \leq C \|\partial_\theta w(t, \cdot)\|_{2,a} \|w(t, \cdot)\|_{2,a} \leq C \|w(t, \cdot)\|_{\mathcal{H}_a^2(S_+^{n-1})}^{\frac{1}{2}} \|w(t, \cdot)\|_{2,a}^{\frac{3}{2}}.$$

Therefore, since $\|w(t, \cdot)\|_{\mathcal{H}_a^2(S_+^{n-1})}$ is bounded, we deduce that $|g(w_B(t))| \leq C e^{3\rho_1^+ t/2}$, which yields the L^∞ -estimate for the JL-supercritical case.

For the JL-critical and JL-subcritical case, the argument is similar, we omit them.

The proof of this lemma is complete.

Lemma 6.5 *Let q be JL-supercritical, then $\xi_1 > 0$, where ξ_1 is the constant given in the previous lemma.*

Proof. Recalling that $\lim_{t \rightarrow +\infty} y_1(t) = \xi_1$, we know that there exists a sequence $\{t_i\}_{i \in \mathbb{N}}$ such that

$$\lim_{i \rightarrow \infty} y_1(t_i) = \xi_1, \quad \lim_{i \rightarrow \infty} y_1'(t_i) = 0.$$

Integrating (49) on (t, t_i) and taking a limit $i \rightarrow \infty$, we obtain

$$-y_1'(t) + (2\rho_1^+ + \sigma)(\xi_1 - y_1(t)) = \int_t^\infty e^{-\rho_1^+ s} f_1(s) ds.$$

Note that both $y_1(t)$ and $y_1'(t)$ converge to zero as $t \rightarrow -\infty$, we deduce that

$$(2\rho_1^+ + \sigma)\xi_1 = \int_{-\infty}^\infty e^{-\rho_1^+ s} f_1(s) ds.$$

By the convexity of the function w^q , we have $-g(w) = (V_B - w)^q - [V_B^q + qV_B^{q-1}(-w)] > 0$ if $w \neq 0$, which yields that $f_1(t) = -g(w_B(t))e_{1B} > 0$ if $w \neq 0$. This and the fact that $2\rho_1^+ + \sigma > 0$, we conclude that $\xi_1 > 0$.

We complete the proof of this lemma.

Lemma 6.6 *Let q be JL-critical, then $\xi_1 > 0$.*

Proof. Since (50) can be written as $(t^2 y_1')' = t e^{-\rho_1 t} f_1$, integrating on $(1, t)$, we obtain

$$t y_1'(t) = t^{-1} y_1'(0) + t^{-1} \int_1^t e^{-\rho_1 s} f_1(s) ds, \quad t > 1,$$

which gives that $\lim_{t \rightarrow \infty} t y_1'(t) = 0$. As a consequence, since $\lim_{t \rightarrow +\infty} y_1(t) = \xi_1$, integrating (50) on (t, ∞) , we obtain

$$-(t y_1'(t) + y_1(t)) + \xi_1 = \int_t^\infty e^{-\rho_1 s} f_1(s) ds.$$

Since $z_1(t) = t e^{\rho_1 t} y_1(t)$, we have $t y_1'(t) + y_1(t) = (-\rho_1 z_1(t) + z_1'(t)) e^{-\rho_1 t}$. Taking $t \rightarrow 0$, we have

$$\xi_1 = (-\rho_1 z_1(0) + z_1'(0)) + \int_0^\infty e^{-\rho_1 s} f_1(s) ds. \quad (54)$$

Similarly, integrating (50) on $(-\infty, t)$, we obtain

$$t y_1'(t) + y_1(t) = \int_{-\infty}^t e^{-\rho_1 s} f_1(s) ds,$$

where we have used the facts that $\lim_{t \rightarrow -\infty} t y_1'(t) = 0$ and $\lim_{t \rightarrow -\infty} y_1(t) = 0$. Since $t y_1'(t) + y_1(t) = (-\rho_1 z_1(t) + z_1'(t)) e^{-\rho_1 t}$, taking $t \rightarrow 0$, we have

$$(-\rho_1 z_1(0) + z_1'(0)) = \int_{-\infty}^0 e^{-\rho_1 s} f_1(s) ds. \quad (55)$$

Add (54) to (55), we conclude that

$$\xi_1 = \int_{-\infty}^\infty e^{-\rho_1 s} f_1(s) ds,$$

which yields $\xi_1 > 0$. The proof of this lemma is complete.

For the JL-subcritical case, if $(\xi_1, \xi_2) = 0$, where (ξ_1, ξ_2) is given in (48), then we will establish the following expansion with a smaller error.

Lemma 6.7 *Let q be JL-subcritical and $(\xi_1, \xi_2) = 0$, then there exists $\xi < 0$ and $C, \varepsilon > 0$ such that*

$$\|e^{-\rho_2^- t} w(t, \theta) - \xi e_2\|_{2,a} \leq C e^{-\varepsilon t}.$$

Moreover the norm $\|\cdot\|_{2,a}$ can be replaced by the norm $\|\cdot\|_\infty$.

Proof. Since $(\xi_1, \xi_2) = 0$, from (47) and (52), we have $|z_1(t)| \leq C e^{3\rho_2^- t/2}$. Then, by the same argument as in the proof of Lemma 6.3, we obtain

$$\sum_{i=3}^{\infty} |z_i(t)| \leq C[e^{\rho_3^- t} + e^{3\rho_2^- t/2}], \quad |z_2(t)| \leq C e^{\rho_2^- t}. \quad (56)$$

Now we set $y_2(t) = e^{-\rho_2^- t} z_2(t)$. Then $y_2(t)$ solves

$$y_2'' + (2\rho_2^- + \sigma)y_2' = e^{-\rho_2^- t} f_2. \quad (57)$$

By (56) we know that $y_2(t)$ is bounded, so there exists a sequence $\{t_i\}_{i \in \mathbb{N}}$ such that $\lim_{i \rightarrow \infty} y_2'(t_i) = 0$. Then we have

$$-e^{(2\rho_2^- + \sigma)t} y_2'(t) = \lim_{i \rightarrow \infty} e^{(2\rho_2^- + \sigma)t_i} y_2'(t_i) - e^{(2\rho_2^- + \sigma)t} y_2'(t) = \int_t^\infty e^{(\rho_2^- + \sigma)s} f_2(s) ds.$$

Hence from (56), we deduce $|y_2'(t)| \leq C e^{\rho_2^- t/2}$. As a consequence, there exists $\xi \in \mathbb{R}$ such that $\lim_{t \rightarrow \infty} y_2(t) = \xi$ and

$$\begin{aligned} \|e^{-\rho_2^- t} w(t, \cdot) - \xi e_2\|_{2,a} &\leq \|e^{-\rho_2^- t} w(t, \cdot) - e^{-\rho_2^- t} z_2(t) e_2\|_{2,a} + \|(e^{-\rho_2^- t} z_2(t) - \xi) e_2\|_{2,a} \\ &\leq \|e^{-\rho_2^- t} [w(t, \cdot) - z_2(t) e_2]\|_{2,a} + \|(y_2(t) - \xi) e_2\|_{2,a} \\ &\leq C e^{-\varepsilon t}, \end{aligned}$$

where in the last inequality we have used (56), the facts that $|z_1(t)| \leq C e^{3\rho_2^- t/2}$ and exponent decay of $y_2'(t)$. Integrating (57) on $(-\infty, \infty)$, we obtain

$$(2\rho_2^- + \sigma)\xi = \int_{-\infty}^{\infty} e^{-\rho_2^- s} f_2(s) ds.$$

since $2\rho_2^- + \sigma < 0$ and $f_2(s) = -g(w_B(t))e_{2B} > 0$, we deduce that $\xi < 0$. Applying the same argument as given in Lemma 6.4, we obtain the L^∞ -estimate.

We complete the proof of this lemma.

Proof of Proposition 2.1 By Lemmas 6.4-6.7, we immediately obtain Proposition 2.1.

Acknowledgment. The author is partially supported by NSFC, No 11101134 and NSFC, No 11371128.

References

- [1] L. Caffarelli and L. Silverstre, *An extension problem related to the fractional Laplacian*, Comm. Partial Differential Equations, 32(2007), 1245-1260.
- [2] M. Chipot, M. Chlebik and M. Fila, *On the solutions to some elliptic equations with nonlinear boundary conditions*, Adv. in Differential Equations 1(1996), 91-110.
- [3] M. Chipot, M. Chlebik, M. Fila and I. Shafrir, *Existence of Positive Solutions of a Semilinear Elliptic Equation in R_+^n with a Nonlinear Boundary Condition*, Journal of Mathematical Analysis and Applications 223, (1998) 429-471.
- [4] J. Dávila, L. Dupaigne and M. Montenegro, *The extremal solution of a boundary reaction problem*, Commun. Pure Appl. Anal. 7, (2008) 795-817.
- [5] Z. Du, Y. Fang and C. Gui, *Positive solutions of a degenerate elliptic equation in the upper half space with a nonlinear boundary condition*, Submitted.
- [6] Z. Du, *Singular solutions of semilinear elliptic equations with fractional Laplacian in entire space*, Communications in Contemporary Mathematics, DOI: 10.1142/S02191997155002484.
- [7] E. B. Fabes, C. E. Kenig and R. P. Serapioni, *The local regularity of solutions of degenerate elliptic equations*, Comm. Partial Differential Equations, 7(1982): 77-116.
- [8] D. Gilbarg and N. S. Trudinger, *Elliptic partial differential equations of second order*, Springer, Berlin, 1998.
- [9] C. Gui, W.-M. Ni and X. Wang, *On the stability and instability of positive steady states of a semilinear heat equation in R^n* , Comm Pure Appl. Math., 45(1992): 1153C1181.
- [10] P. Hajlasz, *Sobolev spaces on an arbitrary metric space*, Potential Analysis, 5(1996), 403-415.
- [11] J. Harada, *Positive solution to the Laplace equation with nonlinear boundary conditions on the half space*, Calc. Var. Partial Differential Equations, 50(1-2)(2014), 399-435.
- [12] B. Hu, *Non existence of a positive solution of the Laplace equation with a nonlinear boundary condition*, Differential Integral Equations, 7(1994): 301-313.
- [13] Y. Li, *Asymptotic behavior of positive solutions of equation $\Delta u + K(x)u^p = 0$ in R^n* , J. Diff. Eqns. 95(1992), 304-330

- [14] Y. Y. Li and M. Zhu, *Uniqueness theorems through the method of moving spheres*, Duke Math. J. 80(1995), 383-417.
- [15] P. Quittner and W. Reichel, *Very weak solutions to elliptic equations with nonlinear Neumann boundary conditions*, Cal. Var. Partial Differential Equations. 32(2008), no. 4, 429-452.